

# Genetic variation in TERT modifies the risk of hepatocellular carcinoma in alcohol-related cirrhosis: results from a genome-wide case-control study

## Info

PUBLIZIERT AM      JOURNAL  
2023-01-05      Gut

## LINK

<https://doi.org/10.1136/gutjnl-2022-327196>

## Autoren

Stephan Buch   Hamish Innes   Philipp Ludwig Lutz  
Hans Dieter Nischalke   Jens U Marquardt  
Janett Fischer   Karl Heinz Weiss   Jonas Rosendahl  
Astrid Marot   Marcin Krawczyk   Markus Casper  
**Frank Lammert**   Florian Eyer   Arndt Vogel  
Silke Marhenke   Johann von Felden   Rohini Sharma  
Stephen Rahul Atkinson   Andrew McQuillin  
Jacob Nattermann   **Clemens Schafmayer**  
Andre Franke   Christian Strassburg  
Marcella Rietschel   Heidi Altmann   Stefan Sulk  
Veera Raghavan Thangapandi   **Mario Brosch**  
Carolin Lackner   Rudolf E Stauber   **Ali Canbay**  
Alexander Link   Thomas Reiberger  
Mattias Mandorfer   Georg Semmler  
Bernhard Scheiner   Christian Datz   Stefano Romeo  
Stefano Ginanni Corradini   William Lucien Irving  
Joanne R Morling   Indra Neil Guha   Eleanor Barnes  
M Azim Ansari   Jocelyn Quistrebert   Luca Valenti  
Sascha A Müller   Marsha Yvonne Morgan  
Jean-François Dufour   **Jonel Trebicka**   **Thomas Berg**  
Pierre Deltenre   Sebastian Mueller   **Jochen Hampe**  
Felix Stickel

## Abstract

**Objective** Hepatocellular carcinoma (HCC) often develops in patients with alcohol-related cirrhosis at an annual risk of up to 2.5%. Some host genetic risk factors have been identified but do not account for the majority of the variance in occurrence. This study aimed to identify novel susceptibility loci for the development of HCC in people with alcohol related cirrhosis.

### Design

Patients with alcohol-related cirrhosis and HCC (cases: n=1214) and controls without HCC (n=1866), recruited from Germany, Austria, Switzerland, Italy and the UK, were included in a two-stage genome-wide association study using a case-control design. A validation cohort of 1520 people misusing

alcohol but with no evidence of liver disease was included to control for possible association effects with alcohol misuse. Genotyping was performed using the InfiniumGlobal Screening Array (V.24v2, Illumina) and the OmniExpress Array (V.24v1-0a, Illumina).

## Results

Associations with variants rs738409 in PNPLA3 and rs58542926 in TM6SF2 previously associated with an increased risk of HCC in patients with alcohol-related cirrhosis were confirmed at genome-wide significance. A novel locus rs2242652(A) in TERT (telomerase reverse transcriptase) was also associated with a decreased risk of HCC, in the combined meta-analysis, at genome-wide significance ( $p=6.41 \times 10^{-9}$ , OR=0.61 (95% CI 0.52 to 0.70)). This protective association remained significant after correction for sex, age, body mass index and type 2 diabetes ( $p=7.94 \times 10^{-5}$ , OR=0.63 (95% CI 0.50 to 0.79)). Carriage of rs2242652(A) in TERT was associated with an increased leucocyte telomere length ( $p=2.12 \times 10^{-44}$ ).

## Conclusion

This study identifies rs2242652 in TERT as a novel protective factor for HCC in patients with alcohol-related cirrhosis.